

On the Geometry of Planar Domain Walls

F. M. Paiva^{*} and Anzhong Wang[†]

Departamento de Astrofísica, Observatório Nacional – CNPq,

Rua General José Cristino 77, 20921-400 Rio de Janeiro – RJ, Brazil

Abstract

The Geometry of planar domain walls is studied. It is argued that the planar walls indeed have plane symmetry. In the Minkowski coordinates the walls are mapped into revolution paraboloids.

PACS numbers: 04.20.Jb, 04.20.Gz, 98.80.Cq, 11.27.+d.

^{*}Internet: fmpaiva@on.br

[†]Internet: wang@on.br

Domain walls are objects formed in the early stages of the evolution of the Universe [1] and have been studied intensively in the past decade or so, mainly because of their notable implications to Cosmology [2]. The first analytic solution of a domain wall with plane symmetry¹ was found by Vilenkin in 1983 [6]. Since then, this solution has been studied by several authors, among them are Gibbons [7] and Wang and Letelier [8], mainly concerning the global structure of the spacetime. Despite of its simplicity, the solution exhibits a very rich global structure. In particular, in each of the three spatial directions there is a horizon. In the direction perpendicular to the wall the horizon is not stable against the perturbations of null fluids [8] and massless scalar fields [9]. In 1984, on the other hand, Ipser and Sikivie [10] found all the planar domain wall solutions that connect two flat regions.

In the existing literature, it is usually believed that those planar domain walls do not really have plane symmetry but spherical one. This belief is mainly due to the early studies of domain walls [10]. In fact, it was showed that *“In Minkowski coordinates, this planar domain wall is not a plane at all, but rather an accelerated sphere”*. This conclusion was also reached in [11]: *“In each case the wall is bent into a closed surface enveloping the original $z > 0$ side of the wall”*, and used quite recently in [12] to study the gravitational

¹Here we use the definition for plane symmetry originally given by Taub [3]. That is, the symmetric plane has three Killing vectors, two represent the translation symmetry and one represents the rotation. Recently, this definition was generalized to a more general case [4, 5].

radiation of planar domain walls.

However, it is well-known that a plane has no one-to-one mapping to a spherical surface. So, it is very curious to see how a planar domain wall is bent into a bubble. In Ref. [8], by considering the analytic maximum extension of the spacetime, it was argued that the geometry of a planar domain wall is a plane. In this short Comment, we shall stress the same argument but in a different direction, and show that in the Minkowski coordinates a planar domain wall is mapped into a revolution paraboloid, instead of a spherically symmetric bubble.

Before proceeding, we would like first to clarify the following concepts: First, when we talk about the geometry of a wall, we mean the geometry of the space-like two-surface of the wall. Second, the Nambu action for a domain wall [2] is defined in a (2+1)-dimensional hypersurface (or a tube), which represents the whole history of the evolution of the wall. Bearing the above in mind, let us consider the Vilenkin domain wall solution

$$ds^2 = e^{-k|z|} \{ dt^2 - dz^2 - e^{kt} (dx^2 + dy^2) \}, \quad (1)$$

where k is a positive constant, and the range of the coordinates is $-\infty < t, z, x, y < +\infty$. The coordinates will be numbered as $\{x^\mu\} = \{t, z, x, y\}$, ($\mu = 0, 1, 2, 3$). The corresponding energy-momentum tensor is given by $T_\nu^\mu = 2k\{1, 0, -1, -1\}\delta(z)$. Thus, it represents a planar domain wall with support only on the hypersurface $z = 0$. The wall has the plane symmetry charac-

terized by the three Killing vectors ∂_x, ∂_y , and $y\partial_x - x\partial_y$, which act on the two-dimensional surfaces t, z constant.

Using Cartan scalar techniques [13], we were able to show that besides these three Killing vectors the spacetime has other three. Indeed, the non-zero Cartan scalars² for the metric (1) are

$$\begin{aligned}\Phi_{11'} &= -2\Phi_{00'} = -2\Phi_{22'} = -\Lambda &= -\frac{1}{4}k^2\delta(z), \\ \nabla\Phi_{11'} &= -\nabla\Phi_{22'} = 3\nabla\Phi_{33'} = -3\nabla\Phi_{00'} &= \\ -\frac{2}{3}\nabla\lambda_{00'} &= \frac{2}{3}\nabla\lambda_{11'} &= -\frac{\sqrt{2}}{12}k^3\delta'(z),\end{aligned}\tag{2}$$

where $\delta(z)$ denotes the Dirac delta function and $\delta'(z)$ its derivative. Outside the hypersurface $z = 0$, all Cartan scalars vanish, therefore, in this region the spacetime is locally flat. Otherwise, from this set of Cartan scalars, one finds that the metric is Segre type $[(1,11)1]$ and its isotropy group is $\text{SO}(2,1)$, which is three dimensional, leading to three Killing vectors (one of them is the spatial rotation $y\partial_x - x\partial_y$ on the plane of symmetry). Since z is the only coordinate appearing on the Cartan scalars, $z = \text{Const.}$ are three dimensional homogeneous hypersurfaces, leading to other three Killing vectors (two of them are the translations ∂_x and ∂_y on the plane of symmetry). Thus, the spacetime has six Killing vectors which act on the $(2+1)$ -hypersurfaces

² The Cartan scalars are basically the components of the Riemann tensor and its covariant derivatives calculated in a constant frame providing a complete local characterization of spacetimes [14, 15]. Spinor components are used and the relevant objects here are the Ricci spinor $\Phi_{AB'}$, the curvature scalar λ and its first symmetrized covariant derivatives $\nabla\Phi_{AB'}$ and $\nabla\lambda_{AB'}$. SHEEP and CLASSI [16, 17] were used in the calculations.

z constant. Three of them, ∂_x , ∂_y and $x\partial_y - y\partial_x$ act on the 2-dimensional surfaces t, z constant. Because these $(2+1)$ -hypersurfaces are time-like, a time-like Killing vector must exist. Thus, the spacetime is at least stationary. In fact, it is locally static, as can be seen by performing the following coordinate transformations

$$\begin{aligned} t &= \bar{t} + \alpha \ln \left[\frac{(\alpha^2 - \rho^2)^{1/2}}{\alpha^2} \right], \\ x &= \frac{\alpha^2 \rho e^{-\bar{t}/\alpha}}{(\alpha^2 - \rho^2)^{1/2}} \cos \phi, \\ y &= \frac{\alpha^2 \rho e^{-\bar{t}/\alpha}}{(\alpha^2 - \rho^2)^{1/2}} \sin \phi, \end{aligned} \tag{3}$$

where $\alpha \equiv 2/k$. In terms of $\bar{t}, \rho$ and ϕ , the metric (1) takes the form

$$ds^2 = e^{-k|z|} \{ f(\rho) d\bar{t}^2 - f^{-1}(\rho) d\rho^2 - \rho^2 d\phi^2 - dz^2 \}, \tag{4}$$

where

$$f(\rho) = 1 - \frac{\rho^2}{\alpha^2}, \tag{5}$$

and $-\infty < \bar{t}, z < +\infty$, $0 \leq \rho < \alpha$ and $0 \leq \phi \leq 2\pi$, which covers part of the spacetime (1). It should be stressed that although the solution is static in the above domain, globally it is not, as Gibbons pointed out in a different way [7].

To study the geometry of the wall, following Ref. [10] (see also Ref. [7]),

let us make the following coordinate transformations

$$\begin{aligned} T &= \frac{e^{-kz/2}}{4k} \left\{ k^2(x^2 + y^2)e^{-kz/2} + 8 \sinh\left(\frac{kt}{2}\right) \right\}, \\ Z &= \frac{e^{-kz/2}}{4k} \left\{ k^2(x^2 + y^2)e^{-kz/2} - 8 \cosh\left(\frac{kt}{2}\right) \right\}, \\ X &= e^{k(t-z)/2}x, \quad Y = e^{k(t-z)/2}y, \end{aligned} \quad (6)$$

or inversely

$$\begin{aligned} t &= \frac{2}{k} \ln \left[\frac{T-Z}{(R^2-T^2)^{1/2}} \right], \quad z = \frac{2}{k} \ln \left[\frac{2}{k(R^2-T^2)^{1/2}} \right], \\ x &= \frac{2X}{k(T-Z)}, \quad y = \frac{2Y}{k(T-Z)}, \end{aligned} \quad (7)$$

in the region $z \geq 0$, where $R^2 \equiv X^2 + Y^2 + Z^2$. Replacing z by $-z$ in the above equations, we will get the coordinate transformations in the region $z \leq 0$. Because of the reflection symmetry, without loss of generality, we shall focus our attention in the region $z \geq 0$.

In terms of T, X, Y , and Z , the metric (1) becomes $ds^2 = dT^2 - dX^2 - dY^2 - dZ^2$ in the region $z \geq 0$, which has the Minkowski form. Thus, we shall refer the coordinates T, X, Y , and Z to as the Minkowski coordinates. From Eq.(6), on the other hand, we find that the two-dimensional surfaces $t = \text{Const.}$, say, $t = t_0$, and $z = 0$ are given by

$$X^2 + Y^2 + Z^2 - T^2 = \frac{4}{k^2}, \quad (8)$$

$$T - Z = \frac{2}{k} e^{t_0 k/2}. \quad (9)$$

Thus, we have

$$X^2 + Y^2 = \frac{4e^{t_0 k/2}}{k} \left\{ \frac{2}{k} \cosh \left(\frac{t_0 k}{2} \right) + Z \right\}, \quad (10)$$

which represents a revolution paraboloid. Therefore, the planar wall looks like a revolution paraboloid in the Minkowski coordinates, instead of a spherically symmetric bubble [10, 11]. Because of the reflection symmetry, we will have the same conclusion when we are working in the region $z \leq 0$.

On the other hand, that the planar wall is not a bubble can be also stressed by considering the Killing vectors. Let us first assume that the wall is a spherically symmetric bubble in the Minkowski coordinates. Then, we know that in this coordinate system the three Killing vectors that define the spherical symmetry are

$$\begin{aligned} \xi_{(1)} &= Y \frac{\partial}{\partial X} - X \frac{\partial}{\partial Y} = y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y}, \\ \xi_{(2)} &= Z \frac{\partial}{\partial Y} - Y \frac{\partial}{\partial Z} \\ &= \frac{e^{-kt}}{4k} \left\{ 4kye^{kt} \frac{\partial}{\partial t} - 2k^2 xy e^{kt} \frac{\partial}{\partial x} - [k^2(y^2 - x^2)e^{kt} + 4(1 + e^{kt})] \frac{\partial}{\partial y} \right\}, \\ \xi_{(3)} &= Z \frac{\partial}{\partial X} - X \frac{\partial}{\partial Z} \\ &= \frac{e^{-kt}}{4k} \left\{ 4kxe^{kt} \frac{\partial}{\partial t} - 2k^2 xy e^{kt} \frac{\partial}{\partial y} - [k^2(x^2 - y^2)e^{kt} + 4(1 + e^{kt})] \frac{\partial}{\partial x} \right\}. \end{aligned}$$

From the above expressions it is clear that only the Killing vector $\xi_{(1)}$ acts on the 2-dimensional space-like surface, $t = \text{Const.}$ and $z = 0$, of the wall, and

$\xi_{(2)}$ and $\xi_{(3)}$ act outside this surface. This contradicts to our assumption, since if the wall has spherically symmetry, then the above three Killing vectors should act on it. Therefore, it is concluded that the geometry of the Vilenkin planar domain wall is plane, on which the three Killing vectors ∂_x, ∂_y , and $y\partial_x - x\partial_y$ act.

All the solutions, which represent infinitely thin domain walls connecting two flat regions, were given in Ref. [10]. By a similar consideration, one can show that all these walls have the same topological structure as the Vilenkin wall. Moreover, this is also true for the Goetz domain wall with non-zero thickness [20, 21].

Recently, Cvetič and co-workers [22] carried out a detailed study of the topology of domain walls, including the planar ones. In particular, it was shown that the only geodesically complete $(2+1)$ -dimensional spacetime $z = \text{Const.}$ is the one whose two-dimensional spatial surfaces, $t, z = \text{Const.}$ have positive curvature, which means that the geometry of the wall in that case is a compact bubble. On the other hand, from the Vilenkin solution (1) we can see that the hypersurfaces $z = \text{Const.}$ are the $(2+1)$ -dimensional de Sitter space written in a coordinate system in which it is geodesically *incomplete*, and that, as a result, the Vilenkin planar domain wall looks like a revolution paraboloid in the four-dimensional Minkowski space rather than a bubble.

In Ref. [12], using the conclusions obtained in Refs. [10] and [11], that all planar domain walls are actually bubbles, the gravitational radiation of a wall was studied. In particular, it was found that to the first-order approximation

the wall does not emit gravitational waves. As we know, it is much easier for an object with plane symmetry to emit gravitational waves than for one with spherical symmetry. So, it would be very interesting to consider the problem directly in the coordinates of Eq.(1). It is most likely that the situation will be different and the final results will support the earlier speculations of Vachaspati, Everett and Vilenkin [23].

Acknowledgment

The authors gratefully acknowledge financial assistance from CNPq.

References

- [1] T.W.B. Kibble, J. Phys. A9, 1387 (1976).
- [2] A. Vilenkin, Phys. Rep. 121, 263 (1985); Phys. Rev. Lett. 72, 3137 (1994) and references therein.
- [3] A.H. Taub, Ann. Math. 53, 472 (1951).
- [4] J. Centrella and R.A. Matzner, Astrophys. J. 230, 311 (1979); Phys. Rev. D25, 930 (1982).
- [5] A.Z. Wang, Phys. Rev. D45, 3534 (1992); P.S. Letelier and A.Z. Wang, Phys. Lett. A182, 220 (1993).
- [6] A. Vilenkin, Phys. Lett. B133, 177 (1983).

- [7] G.W. Gibbons, Nucl. Phys. B394, 3 (1993).
- [8] A.Z. Wang and P.S. Letelier, “*Instability of Cosmological Event and Cauchy Horizons in the Spacetime with Plane Symmetry*,” CNPq-ON-preprint (1995).
- [9] A.Z. Wang, Phys. Lett. B277, 49 (1992).
- [10] J. Ipser and P. Sikivie, Phys. Rev. D30, 712 (1984).
- [11] J. Ipser, Phys. Rev. D30, 2452 (1984); 36, 1933 (1987).
- [12] H. Kodama, H. Ishihara, and Y. Fujiwara, Phys. Rev. D50, 7292 (1994).
- [13] M.A.H. MacCallum and J.E.F. Skea, “SHEEP: *A Computer Algebra System for General Relativity*”, in “Algebraic Computing in General Relativity: Lecture Notes from the First Brazilian School on Computer Algebra”, Vol. 2, edited by M. J. Rebouças and W. L. Roque (Oxford University Press, Oxford, 1994).
- [14] É. Cartan, “*Leçons sur la Géométrie des Espaces de Riemann*” (Gauthier-Villars, Paris, 1951).
- [15] A. Karlhede, Gen. Rel. Grav. 12, 693 (1980).
- [16] I. Frick, “SHEEP *Users Guide*”, Institute of Theoretical Physics, University of Stockholm Report 77–14 (1977).

- [17] J.E. Åman, “*Manual for CLASSI - Classification Programs for Geometries in General Relativity (Third Provisional Edition)*”, University of Stockholm Report (1987).
- [18] S.W. Hawking and G.F.R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, Cambridge, England, 1973).
- [19] M. Cvetič, R.L. Davis, S. Griffies, and H.H. Soleng, Phys. Rev. Lett. 70, 1191 (1993); M. Cvetič, S. Griffies, and H.H. Soleng, Phys. Rev. D48, 2613 (1993).
- [20] G. Goetz, J. Math. Phys. 31, 2683 (1990).
- [21] A.Z. Wang and P.S. Letelier, “*Local and Global Structure of a Thick Domain Wall Spacetime*”, CNPq-ON-preprint (1994).
- [22] M. Cvetič, S. Griffies, and H.H. Soleng, Phys. Rev. Lett. 71, 670 (1993); Phys. Rev. D48, 2613 (1993).
- [23] T. Vachaspati, A.E. Everett, and A. Vilenkin, Phys. Rev. D30, 2046 (1984).